CLAIMS

1. A polynucleotide of any one of (a) to (d):

5

10

15

20

30

- (a) a polynucleotide encoding a polypeptide comprising an amino acid sequence of any one of SEQ ID NO: 2, 4, 6, or 8;
- (b) a polynucleotide comprising the coding region of a nucleotide sequence of any one of SEQ ID NO: 1, 3, 5, or 7;
- (c) a polynucleotide encoding a polypeptide having an activity of causing a keratinocyte to differentiate into a stratified epithelial cell when expressed in the keratinocyte, wherein the polypeptide comprises an amino acid sequence with a substitution, deletion, insertion, and/or addition of one or more amino acids in the amino acid sequence of any one of SEQ ID NO: 2, 4, 6, or 8; and
- (d) a polynucleotide encoding a polypeptide having an activity of causing a keratinocyte to differentiate into a stratified epithelial cell when expressed in the keratinocyte, wherein the polynucleotide hybridizes under stringent conditions with a DNA comprising a nucleotide sequence of any one of SEQ ID NO: 1, 3, 5, or 7.
- 2. The polynucleotide of claim 1 that is a gene involved in keratinocyte differentiation or proliferation, wherein the polynucleotide encodes a secreted protein.
- 3. A polypeptide encoded by the polynucleotide of claim 1 or 2.
- 4. A vector into which the polynucleotide of claim 1 or 2 is inserted.
- 5. A host cell carrying the polynucleotide of claim 1 or 2 or the vector of claim 4.
 - 6. A method for producing the polypeptide of claim 3, comprising the steps of culturing the host cell of claim 5, and recovering a produced polypeptide from the host cell or its culture supernatant.
 - 7. A gene complex related to keratinocyte differentiation or proliferation, comprising each of (1) a Kdap gene; (2) a dermokine-α gene; (3) a dermokine-β gene; and (4) a suprabasin gene, wherein regulation of gene expression is commonly controlled.
- 8. A polynucleotide that hybridizes specifically with the polynucleotide of claim 1 or 2, wherein the polynucleotide has a chain length of at least 15 nucleotides.

- 9. An antisense polynucleotide against the polynucleotide of claim 1 or 2, or a portion thereof.
- 10. An antibody that binds to the polypeptide of claim 3.

10

15

20

25

- 11. An agent for inducing keratinocyte differentiation, comprising a dermokine-α protein or a dermokine-β protein as an active ingredient.
 - 12. An inhibitor of keratinocyte differentiation, comprising a compound selected from the group of (a) to (d):
 - (a) an antisense nucleic acid against a transcript of a dermokine-α gene or a dermokine-β gene;
 - (b) a nucleic acid having a ribozyme activity that specifically cleaves a transcript of a dermokine- α gene or a dermokine- β gene;
 - (c) a nucleic acid having an effect of inhibiting the expression of a dermokine- α gene or a dermokine- β gene through an RNAi effect; and
 - (d) an antibody that binds to a dermokine- α protein or a dermokine- β protein.
 - 13. A method of screening for an inducer of keratinocyte differentiation, comprising steps (a) to (c):
 - (a) contacting a cell that expresses a dermokine-α protein or a dermokine-β protein with a test compound;
 - (b) measuring an expression level or an activity of a dermokine- α protein or a dermokine- β protein in the cell; and
 - (c) selecting a compound that increases the above expression level or activity, compared to in the absence of contact with the test compound.
 - 14. A method of screening for an inhibitor of keratinocyte differentiation, comprising steps (a) to (c):
 - (a) contacting a cell that expresses a dermokine-α protein or a dermokine-β protein with a test compound;
- (b) measuring an expression level or an activity of a dermokine- α protein or a dermokine- β protein in the cell; and
 - (c) selecting a compound that decreases the above expression level or activity, compared to in the absence of contact with the test compound.
- 15. A method of screening for an inducer of keratinocyte differentiation, comprising steps (a) to (c):

- (a) coexisting a test compound and a keratinocyte with a dermokine- α protein, a dermokine- β protein, or a cell expressing these proteins;
 - (b) measuring the differentiation of a keratinocyte into a stratified epithelial cell; and
- (c) selecting a compound that increases differentiation into a stratified epithelial cell, compared to in the absence of the test compound.

5

10

25

35

- 16. A method of screening for an inhibitor of keratinocyte differentiation, comprising steps (a) to (c):
- (a) coexisting a test compound and a keratinocyte with a dermokine- α protein, a dermokine- β protein, or a cell expressing these proteins;
 - (b) measuring the differentiation of a keratinocyte into a stratified epithelial cell; and
- (c) selecting a compound that decreases differentiation into a stratified epithelial cell, compared to in the absence of the test compound.
- 17. A method for examining whether or not a subject cell is a cancer cell derived from a stratified epithelium, comprising steps (a) and (b):
 - (a) measuring an expression level or an activity of a dermokine- α protein or a dermokine- β protein in a subject cell; and
- (b) determining that the subject cell is a cancer cell derived from a stratified epithelium when the above expression level or activity is different from a control.
 - 18. A method for diagnosing a squamous epithelial cancer or a basal cell cancer in a subject, comprising steps (a) and (b):
 - (a) determining whether or not a cell sample prepared from a subject is a cancer cell derived from a stratified epithelium, using the method of claim 17; and
 - (b) determining that the subject is affected with a squamous epithelial cancer or a basal cell cancer when the cell sample is determined to be a cancer cell derived from a stratified epithelium in the above step.
- 19. A method for diagnosing a skin disease in a subject, comprising steps (a) and (b):
 - (a) measuring an expression level or activity of a dermokine- α protein or a dermokine- β protein in a test sample prepared from a subject; and
 - (b) determining that the subject is affected with a skin disease when the above expression level or activity is different from a control.
 - 20. The method for diagnosing of claim 19, wherein the skin disease is a xeroderma, psoriasis, or

ichthyosis.